



Arthur Schlyer, M.D.
South Tampa, Westchase & NPR
Phone 727-847-1825
Fax 727-849-4855

Priapus Shot[®]

- ❖ Please do not use aspirin or other NSAID pain relievers (Advil, Motrin, Aleve) for one week after the procedure. You may use acetaminophen (Tylenol) as directed on the label and/or apply an ice pack to the area for any discomfort.
- ❖ Arnica pills or gel can be purchased at homeopathic/vitamin stores, if desired. Arnica is an herb thought to help reduce any bruising. Please follow any manufacturers instructions on the bottle and discontinue if any rash or discomfort develops.
- ❖ Results of this procedure are usually seen within a few hours and continue to improve for up to 18 weeks. You should notice increased blood flow to the area with increased sensation and sensitivity.

Priapus Shot Pump Protocol

The doctor will instruct you on the use of the pump. It should be used for 10 minutes at a pressure of 5 to 10 mmHg immediately after the shot.

The pump should be used twice a day at a pressure of 5 to 10 mmHg for 10 to 15 minutes for at least 6 weeks. These two sessions can be back-to-back separated by a 5-minute rest. The shot works for erectile dysfunction without the pump. But for maximal growth, the pump is recommended.

Please call our office with any questions or concerns at 727-847-1825.